

A multiplex PCR approach to simultaneously genotype potato towards the resistance alleles *Ry-f_{sto}* and *Ns*

Kamil Witek · Danuta Strzelczyk-Zyta ·
Jacek Hennig · Waldemar Marczewski

Received: 18 April 2006 / Accepted: 3 May 2006 / Published online: 28 September 2006
© Springer Science+Business Media B.V. 2006

Abstract A simple and robust multiplex PCR approach was developed for detection of the alleles *Ry-f_{sto}* and *Ns* conferring resistance of potato to *Potato Virus Y* (PVY) and *Potato Virus S* (PVS), respectively. Cleaved amplified polymorphic sequence (CAPS) markers GP122₅₆₄ linked to *Ry-f_{sto}* and SC811₂₆₀ linked to *Ns* were amplified in one PCR reaction and identified after simultaneous digestion of the amplicons with restriction enzymes *EcoRV* and *MboI*. Effectiveness of this procedure for marker-assisted selection was confirmed in 55 potato cultivars.

Keywords MAS · Multiplex PCR · Potato · PVS · PVY · Resistance

Potato Virus Y (PVY) and *Potato Virus S* (PVS) are pathogens that occur worldwide in potato (*Solanum tuberosum* ssp. *tuberosum*) crops (Brunt 2001). Breeding of resistant potato cultivars is the most effective and environmentally safe strategy to achieve protection against Potato

viruses (Świeżyński 1994). From the 1930s to the present wild *Solanum* species have been widely used as sources of resistance genes in cultivated potato breeding programs (Valkonen et al. 1996). The gene *Ns* derived from *Solanum tuberosum* ssp. *andigena* conferred resistance to PVS in potato (Marczewski et al. 1998), whereas *S. stoloniferum* was a donor of the gene *Ry-f_{sto}* responsible for extreme resistance of potato to PVY (Flis et al. 2005). Potato plants possessing *Ry-f_{sto}* and/or *Ns* remained symptomless after mechanical inoculation. *Ns* and *Ry-f_{sto}* have been mapped on potato chromosomes VIII (Marczewski et al. 2002) and XII (Flis et al. 2005), respectively. The development of DNA markers linked to desired traits, including disease resistance genes, becomes more and more helpful for marker-assisted selection in potato breeding programs (Barone 2004). An inter-simple sequence repeat (ISSR) marker UBC811₆₆₀ was applied to identify the allele *Ns* in cultivars resistant to PVS (Marczewski 2001). A cleaved amplified polymorphic sequence (CAPS) marker GP122₇₁₈ was successfully used to detect several Polish and German cultivars expressing extreme resistance to PVY (Flis et al. 2005). GP122₇₁₈ and UBC811₆₆₀ have been cloned and sequenced according to Marczewski et al. (2006). UBC811₆₆₀ (accession DQ415915) did not share significant homology to any sequences deposited in data bases. *EcoRV* recognition site, being diagnostic in

K. Witek · J. Hennig
Institute of Biochemistry and Biophysics, Polish
Academy of Sciences, Pawińskiego 5A,
02106 Warsaw, Poland

D. Strzelczyk-Zyta · W. Marczewski (✉)
Plant Breeding and Acclimatization Institute,
Platanowa 19, 05831 Młochów, Poland
e-mail: w.marczewski@ihar.edu

detection of the allele *Ry-f_{sto}* (Flis et al. 2005), was found for all seven randomly chosen bacterial clones, possessing a 614 bp marker fragment of GP122 (DQ151984-91). In contrast, none of seven 614 bp inserts (DQ151992-97) amplified in a PVY-susceptible plant contained *EcoRV* site.

Here, we report a multiplex PCR approach for detection of the alleles *Ry-f_{sto}* and *Ns*. New PCR primers were designed as previously described (Marczewski et al. 2006) to amplify a 564 bp fragment of GP122 and a 454 bp fragment, designated as SC811₄₅₄, for the *Ns*-linked marker. The PCR amplification was performed in 20 µl of 20 mM Tris-HCl pH 8.4, 50 mM KCl, 2.5 mM MgCl₂, 0.1 mM of each deoxynucleotide, 0.2 µM of each primer (f: 5'-TATTTTAgGGGTACTTCTTTCTTATGTT-3'; r: 5'-CTGTCAAAAAAATTCAGTCCCTTCTAATCCACTAT-3') for GP122 and 0.25 µM of each primer (f: 5'-CGAACAAAATACGTAATGCATTGAATAA-3'; r: 5'-GACCTATATCAGTCCCTTCTAATCCACTAT-3') for SC811₄₅₄, containing 1 U *Taq* DNA Polymerase (Invitrogen, Carlsbad, CA, or Fermentas, St. Leon-Rot,) and 30 ng of genomic DNA as a template. A total genomic DNA was isolated using classical CTAB- method (Weigel and Glazebrook 2002) or as described in Flis et al. (2005). The PCR parameters were: 94°C for 60 s followed by 30 cycles of 93°C for 15 s, 56°C for 20 s, 72°C for 60 s and a final extension time of 5 min at 72°C. Electrophoretic conditions were used as previously described (Flis et al. 2005). The amplicons produced by the multiplex PCR were digested simultaneously with the restriction enzymes *EcoRV* and *MboI*.

Fifty-five potato cultivars bred in Poland and established for resistance level to PVY and PVS were tested. Nineteen PVY-extreme resistant cultivars listed in Flis et al. (2005) and five newly evaluated: Danusia, Jasia, Sonda, Śleza and Ursus, possessed the gene *Ry-f_{sto}*. Among these, Barycz, Klepa, Meduza and Omulew carried both *Ry-f_{sto}* and *Ns*. Out of 23 cvs indicating hypersensitivity or susceptibility to PVY (Flis et al. 2005) only cv. Neptun was known to carry the gene *Ns* (M. Chrzanowska, personal communication). The other 22 and the following eight cvs: Bartek, Czaplą, Ekra, Osa, Tara, Tokaj, Wawrzyn and Zebra are known to lack *Ry-f_{sto}* and *Ns*. The




Fig. 1 Multiplex PCR for detection of CAPS markers GP122₅₆₄ linked to *Ry-f_{sto}* and SC811₂₆₀ linked to *Ns*. Cultivars having *Ry-f_{sto}*: Alicja, Anielka, Beata and Nimfy are shown in lanes 1–4, respectively. Lanes 5–8 show cultivars possessing both *Ry-f_{sto}* and *Ns*: Barycz, Klepa, Omulew and Meduza. Lane 9: cv. Neptun with *Ns*. Lanes 10–12 show cvs Tokaj, Wawrzyn and Zebra, that are known to lack *Ry-f_{sto}* and *Ns*. Lane M contains the 100 bp DNA ladder. GP122₅₆₄ and SC811₂₆₀ are indicated by arrows

presence of the 564 bp *EcoRV* restriction fragment of the GP122 amplicon in all 24 PVY-extreme resistant cultivars indicated that the sequence GP122₅₆₄ linked to *Ry-f_{sto}* did not possess a restriction site for *MboI*. On the other hand, *MboI*-digested SC811₄₅₄ revealed an approximately 260 bp informative restriction fragment, SC811₂₆₀, which was useful for tagging the locus *Ns* in the reference population *Ns* (Marczewski et al. 1998). SC811₂₆₀ was found in all five cultivars resistant to PVS and was not observed in 30 cultivars which lacked *Ry-f_{sto}* and *Ns*. *EcoRV* did not interfere with identifying of SC811₂₆₀. The genotyping of 12 cultivars is shown in Fig. 1.

The genes *Ns* and *Ry-f_{sto}* were introgressed into many European potato gene pools (Flis et al. 2005; Marczewski et al. 1998). Hence, this method provides a tool for the simultaneous tagging of these resistance loci in various breeding programs. To our knowledge it is the first example of the simple and reproducible multiplex PCR that can considerably simplify screening of potato for PVY and PVS resistance through marker-assisted selection.

References

- Barone A (2004) Molecular marker-assisted selection for potato breeding. *Am J Potato Res* 81:111–117
- Brunt AA (2001) Potato Virus M (PVM; Genus *Carlavirus*). In: Loebenstein G, Berger PH, Brunt AA, Lawson RH (eds), *Virus and virus-like diseases of*

- potatoes and production of seed-potatoes. Kluwer, Dordrecht, Boston, London, pp 101–108
- Flis B, Hennig J, Strzelczyk-Zyta D et al. (2005) The *Ry-f_{sto}* gene from *Solanum stoloniferum* for extreme resistant to *Potato Virus Y* maps to potato chromosome XII and is diagnosed by PCR marker GP122₇₁₈ in PVY resistant potato cultivars. *Mol Breed* 15:95–101
- Marczewski W (2001) Inter-simple sequence repeat (ISSR) markers for the *Ns* resistance gene in potato (*Solanum tuberosum* L.). *J Appl Genet* 42:139–144
- Marczewski W, Hennig J, Gebhardt C (2002) The *Potato Virus S* resistance gene *Ns* maps to potato chromosome VIII. *Theor Appl Genet* 105:564–567
- Marczewski W, Ostrowska K, Zimnoch-Guzowska E (1998) Identification of RAPD markers linked to the *Ns* locus in potato. *Plant Breed* 117:88–90
- Marczewski W, Strzelczyk-Zyta D, Hennig J et al. (2006) Potato chromosomes IX and XI carry genes for resistance to Potato Virus M. *Theor Appl Genet* 112:1232–1238
- Świeżyński KM (1994) Inheritance of resistance to viruses. In: Bradshaw JE, Mackay GR (eds), *Potato genetics*. CAB International, Wallingford, pp 339–363
- Valkonen JPT, Jones RAC, Slack SA et al. (1996) Resistance specificities to viruses in potato: standardization of nomenclature. *Plant Breed* 115:433–438
- Weigel D, Glazebrook J (2002) How to isolate a gene defined by a mutation. In: Weigel D, Glazebrook J (eds), *Arabidopsis: a laboratory manual*. CSHL Press, Cold Spring Harbor, New York, pp 143–170